

doi: 10.3325/cmj.2011.52.665

Caries prevalence among schoolchildren in Zagreb, Croatia

Walter Dukić¹, Barbara Delija², Olga Lulić Dukić¹

¹Department of Pediatric Dentistry, School of Dental Medicine, University of Zagreb, Zagreb, Croatia

²Dental Polyclinic Zagreb, Zagreb, Croatia

Aim To investigate the prevalence of dental caries and treatment needs in schoolchildren aged 7-14 years from Zagreb.

Methods Dental examinations based on the World Health Organization criteria were performed on 1168 children in the period 2009-2010. The teeth were clinically examined with standard dental instruments using visual-tactile method under standard dental light. We recorded the clinical indexes of decayed, missed, and filled teeth (DMFT and dmft; upper-case letters refer to permanent and lower-case letters to primary teeth) and decayed, missed, and filled surfaces (DMFS), as well as the significant caries index (SiC).

Results The median DMFT and DMFS of all children were 3 and 4, respectively. The median DMFT and DMFS of 12-year-old children were 4 and 5, respectively. The highest median DMFT score of 7 was found among 14-year-old children. There was a significant difference between age groups (7-10 years and 11-14 years) in DMFT and DMFS. Among 8-year-old children, the median dmft index was the highest (5.5) and SiC index was 7.4. As far as the location of caries on the surface of the first permanent molar is concerned, caries occurred mostly in the central occlusal surface (27.6%).

Conclusion Our results showed a high caries prevalence among schoolchildren in Zagreb, indicating a need for an extensive program of primary oral health care.

Received: August 7, 2011

Accepted: November 25, 2011

Correspondence to:

Walter Dukić
School of Dental Medicine,
University of Zagreb
Gundulićeva 5
Zagreb 10000, Croatia
dukic@sfzg.hr

Dental caries is a public health problem that affects pre-school and high-school children throughout the world, leading to pain, chewing difficulties, speech problems, general health disorders, psychological problems, and lower quality of life (1-5). Although advanced preventive procedures significantly decrease the prevalence of caries in the population, fissure caries on occlusal surfaces and buccal/lingual surfaces is still a considerable problem (6). In permanent dentition, teeth with deep pits and fissures have a higher risk of caries than smooth tooth surfaces (7). Also, caries more frequently affects primary than permanent teeth due to difference in enamel structure, lack of oral hygiene, or absence of preventive measures in oral health (8). Information on caries prevalence and severity represents the basis for caries prevention programs and indicates treatment necessity in the population (9). Measures of caries prevalence are indexes of decayed (D), missed (M), and filled (F) permanent teeth (T) or surfaces (S), ie, DMFT and DMFS indexes, and decayed (d), missed (m), and filled (f) primary teeth (t) or surfaces (s), ie, dmft and dmfs indexes (10). Many reports have indicated that dental caries is decreasing in many European countries and the USA (7,11-21). With a reduction of interproximal smooth surfaces caries, an increased number of cases of occlusal caries inside pits and fissures was recorded (7). The risk of caries significantly increases among adolescents with a high frequency of cariogenic snack consumption. It is also influenced by maternal socio-economic background and educational level (11), as well as dietary, hygienic, and other socio-economic factors, which demonstrates the importance of preventive educational programs and a comprehensive caries prevention scheme for schoolchildren (12).

The aim of this study was to assess the caries prevalence in two primary schools in Zagreb using the indexes for decayed, missed, and filled teeth/surfaces for primary and permanent teeth (DMFT, DMFS, dmft) and significant caries index (SiC).

METHODS

The study was conducted in two primary schools in Zagreb, Croatia, in 2009 and 2010. The two schools were selected because of their location near the city center and proximity to each other (less than 1.5 km). According to the National Census 2001, 52% of residents of this city district had high school education and 27% had college or university education (22-24). A total of 1168 children, 583 male and 585 female, were examined for dental caries.

The study was approved by the Ethics Committee of

the School of Dental Medicine, in Zagreb, Croatia. The procedures and possible discomforts were explained to the children and their parents and an informed consent was obtained before the examination.

Oral examinations were performed by two specialists in pedodontics who had been validated two months before the examination. The inter-examiner reliability had a kappa value of 0.957 based on the examination of 20 children of different ages. After having assessed the completed questionnaires with general information about each child, the examination started with prophylactic cleaning of all teeth surfaces in a dental unit using standard illumination. Prophylactic cleaning was conducted using a slow rotating bristle brush on a micromotor (6000/min), with professional toothpaste Klint (Voco, Cuxhaven, Germany). Teeth were rinsed with water and air, dried, and a dry working field was made using the saliva ejector and cotton rolls to isolate teeth from buccal/labial mucosa and tongue. Examinations for dental caries were conducted using World Health Organization (WHO) criteria and procedures (25). All information was recorded on a specially printed form, including the name, age, address, school, date of examination, and dental record according to the WHO. The status of each tooth was coded using visual-tactile method for analysis. According to the WHO criteria, the dental status of primary teeth was coded with the letters from A to G (healthy tooth, filled, decayed, fissure sealing, etc). Permanent teeth were coded with the numbers from 0-9 according to the WHO codes, with 0 referring to "healthy," 1 referring to "decayed," etc (25). All examined teeth were included in the calculation of the final SiC, DMFT, DMFS, and dmft indexes. The indexes were compared between the age groups, male and female participants, and with the WHO criteria (26,27), which were based on the examination of 12-year-old children. The first molars were thoroughly analyzed regarding the caries localization on tooth surface (mesial, central, or distal molar surface) and WHO criteria.

The indexes were calculated as follows:

DMFS: total number of decayed (D), missing/extracted (M), or filled (F) permanent teeth surfaces (S).

DMFT: total number of decayed (D), missing (M), or filled (F) permanent teeth (T).

DMFT/DMFS: the relative relationship between DMFT and DMFS.

dmft: total number of decayed (d), missing (m) or filled (f) primary teeth (t).

SiC: the mean DMFT for the third of the population with the highest caries scores.

Chi square test was used to study the association between the prevalence of dental caries and sex and age. Kruskal-Wallis and Kolmogorov-Smirnov tests were also used. The Microsoft Office Excel 2007 for Windows (Microsoft Corporation, Redmond, WA, USA) was used for the entry of data on oral health status and for creating the charts. The data were statistically processed using SPSS 11.5 for Windows (SPSS Inc., Chicago, IL, USA).

RESULTS


A total of 1168 children was surveyed, 583 boys and 585 girls, with the mean age of 10.8±2.3 years. Boys and girls did not significantly differ in the number of permanent

teeth at a value of α=0.05 (P=0.141) and in DMFT and DMFS (P>0.05).

DMFT and DMFS increased with age (Figure 1). Children aged 14 had a mean DMFT and DMFS of 7.2 and 10.2, and median of 7 and 8, respectively (web-extra material). Children aged 12 had a mean DMFT and DMFS of 4.8 and 6.9, and median of 4 and 5, respectively. The Kruskal-Wallis test showed that older children (11-14 years) had a significantly higher DMFT (χ²=286.958 df=1 P<0.001) and DMFS (χ²=278.255 df=1, P<0.001) than younger children (7-10 years) (Table 1).

Kolmogorov-Smirnov test showed that both the original and transformed (logarithmic) values of DMFT and DMFS were not normally distributed. The DMFT and DMFS standard deviation had the same or even greater value than the arithmetic mean hence the use of arithmetic mean and standard deviation as descriptive indicators was not justified. Large variability in DMFT and DMFS was additionally evident from the coefficient of quartile deviation (they exceed the value of 0.6). Still, mean values±standard deviations were calculated for comparison of caries indexes with those from similar studies. The mean DMFT for age subgroup 7-10 years was 2.0 and for the age subgroup 11-14 years it was 5.9, while the mean index of SiC for the age

Figure 1.



Median number of decayed, missed, and filled teeth (DMFT) (circles), decayed, missed, and filled surfaces (DMFS) (squares) according to age (N=1168).

TABLE 2. Means and standard deviations of decayed, missed, and filled teeth (DMFT), decayed, missed, and filled surfaces (DMFS) and significant caries (SiC) indexes according to sex and age in schoolchildren in Zagreb

	N	SiC	DMFT	DMFS	DMFT/DMFS × 100
Sex:					
male	583	4.1 ± 3.9	5.8 ± 6.4	82.9	
female	585	4.0 ± 4.0	5.5 ± 6.6	84.0	
Age (years):					
7-10	551	4.0	2.0 ± 1.7	2.5 ± 2.6	88.7
11-14	617	10.8	5.9 ± 4.5	8.4 ± 7.6	79.6
Total	1168	7.4	4.1 ± 4.0	5.6 ± 6.5	83.4

TABLE 1. Decayed, missed, and filled teeth (DMFT) and decayed, missed, and filled surfaces (DMFS) indexes according to sex and age in schoolchildren in Zagreb

	No. (%)	DMFT		DMFS		DMFT/DMFS × 100
		median	coefficient of quartile deviation	median	coefficient of quartile deviation	
Sex:						
male	583 (49.9)	3.00	0.71	4.00	0.78	82.9
female	585 (50.1)	3.00	0.71	4.00	0.75	84.0
Age (years):						
7-10	551 (47.2)	2.00*	1.00	2.00**	1.00	88.7
11-14	617 (52.8)	5.00*	0.60	6.00**	0.60	79.6
Total	1168 (100.0)	3.00	0.71	4.00	0.78	83.4

*Statistical significance (P<0.001).

subgroup 7-10 years was 4.0, and for the age subgroup 11-14 years it was 10.8 (Table 2).

There were 52.4% of first permanent molars with caries on occlusal surface, and only 0.84% in which the caries expanded to complete molar surface (mesial, occlusal, and distal) (Table 3). The caries occurred mostly in the central part of occlusal surface (27.6%) and it was mostly filled (26.9%) with tooth restoration (Table 4).

Missing (extracted) permanent teeth (M) were very rare in this population: of 1168 children, only 51 (4.3%) were missing 1, 2, 3, 4, or 5 permanent teeth, with mean M of 0.08. Filled teeth (F) were more common, with 40.9% of children having at least one filled tooth and a mean F of 1.1. Only


24.0% of children had no caries (D), while 18.1% had one carious tooth, 15.2% had two carious teeth, 10.7% had three carious teeth, and 16.1% had four or more carious teeth. The number of carious teeth (D) increased with age (Figure 2).

There were 54.7% of all children with no caries on primary teeth, 8.7% with a single incident of caries, and 7.7% with two. There were 7.1% of children with one extracted primary tooth, 79.3% had no fillings, 9.8% only had one, 4.6% had two, and 6.3% had three or more.

DISCUSSION

This study showed that our population neither reached the WHO goal for 2010 of DMFT in 12-year-olds below 1, nor

Figure 2.



Average number of permanent teeth (rhombs) and average number of teeth with caries (squares) in children examined according to age (N=1168).

TABLE 4. Detailed analysis of all first molar surfaces according to World Health Organization (WHO) criteria (25)

WHO criteria	No. (%) of teeth with caries on		
	mesial molar surface	distal molar surface	central occlusal surface
Sound	4387 (93.9)	4515 (96.6)	1239 (26.5)
Decayed	134 (2.9)	57 (1.2)	1289 (27.6)
Filled, with decay	20 (0.4)	14 (0.3)	267 (5.7)
Filled, no decay	95 (2.0)	52 (1.1)	992 (21.2)
Missing, as a result of caries	17 (0.4)	17 (0.4)	17 (0.4)
Fissure sealant	0 (0)	0 (0)	852 (18.2)
Unerupted tooth (crown)	15 (0.3)	15 (0.3)	15 (0.3)
Not recorded	4 (0.08)	2 (0.04)	1 (0.02)
Total	4672	4672	4672

TABLE 3. Analysis of caries according to first permanent molar surface in schoolchildren in Zagreb

Variable	No. (%) of children with			Total	χ^2 test
	without caries	partial caries (mesial, central, or distal)	complete caries (mesial + central + distal)		
Sex:					
male	638 (46.4)	723 (52.6)	13 (1.0)	1374	$\chi^2_2 = 0.432$ $P = 0.806$
female	574 (47.1)	634 (52.1)	9 (0.8)	1217	
Age:					
7-10*	734 (51.9)	677 (47.8)	4 (0.3)	1415	$\chi^2_7 = 66.526$ $P < 0.001$
11-14*	478 (40.6)	680 (57.8)	18 (1.5)	1176	
7*	73 (57.9)	53 (42.1)	-	126	
8*	377 (56.3)	292 (43.6)	1 (0.1)	670	
9*	175 (49.7)	177 (50.3)	-	352	
10*	109 (40.8)	155 (58.1)	3 (1.1)	267	
11*	105 (47.3)	117 (52.7)	-	222	
12*	142 (45.8)	161 (51.9)	7 (2.2)	310	
13*	82 (35.3)	143 (61.6)	7 (3)	232	
14*	149 (36.2)	259 (62.9)	4 (0.9)	412	
Total	1212 (46.8)	1357 (52.4)	22 (0.8)	2591	

*Statistical significance ($P < 0.001$).

the one of below 3 for 2000 (26,27). The mean and median DMFT of children aged 12 years in our study was 4.8 and 4, respectively. Epidemiological indexes this high could be explained by the lack of national preventive programs, insufficient number of pediatric dentists in Croatia, and the lack of preventive and educational measures.

There are noticeable differences in the measured indexes between this and other similar studies. In surveys conducted between 1991 and 1995, the mean dmft in primary dentition in children aged 5-7 years ranged between 0.9 and 8.5 (28). It was lowest in Spain (1.0 dmft) and Denmark (1.3 dmft). National mean dmft values below 2.0 were also reported in Finland, the Netherlands, and Norway (28). Ireland had the lowest mean dmft value of 0.9 (28). All of the mentioned countries had a lower mean dmft than this study, which was 5.0 for 7-year-old children.

The mean DMFT in the majority of countries was below 3.0, and in the countries of North-Western Europe and the USA it was below 2.0 (29). However, other European populations, particularly those living in the Mediterranean area, had different results. Twelve-year-old Sicilian schoolchildren had the mean DMFT of 2.88 and their Sardinian peers of 2.4 (11,12). Twelve-year-old Greek children had a DMFT from 2.77 to 6.74, and parental education status, reason for visiting a dentist, and oral hygiene were strong determinants for caries experience (13). Spain had the mean DMFT in 12-year-old children of 1.33, with a goal to reduce it to below 1.0 by 2015 (19,20). Another study from Spain showed the mean DMFT of 2.43 in 12-year-old immigrant children and 0.99 in Spanish children (30). The mean DMFT in German 8-9-year-olds was 0.7 and in Hungary it was 0.4 (21).

On the other hand, 12-year-olds in many countries had a mean DMFT higher than 3.0, such as Latvia (7.7), Poland (5.1), Ukraine (4.4), Hungary (4.3), Lithuania and Belarus (3.8), Russia (3.7), Romania, Portugal, and Bulgaria (28). Also, Bosnia and Herzegovina, Montenegro, and Kosovo reported DMFT mean values above 3 (15-18). These results are similar to our mean DMFT of 4.8 in 12-year-old children. Considering the fact that this study was conducted in 2009-2010, this leads to a conclusion that the situation in Zagreb is not satisfactory.

The SiC index is a reliable tool for studying children with a high risk of caries (31). In Italy (Sardinia), there was a decrease in the DMFT and SiC in the period from 1989 to 2004 (32). The mean DMFT index decreased from 4.3 ± 3.1 in 1989 to 0.8 ± 1.5 in 2004 and the SiC index decreased

from 7.8 (1989) to 3.9 (2004). In our study, the SiC index was extremely high, 7.4. This result indicates a great need for thoroughly planned prevention and restoration of dental caries in Croatia. However, the SiC index obtained in this study is indeed lower than that obtained in a 2008 study conducted in rural parts of Croatia on 301 children aged 3-14 years (10.89) (33). This can indicate that in the past few years a better oral hygiene and sense of oral health has been accomplished. In the same study, the mean DMFT in children aged 11-14 years was 6.7, mean DMFS 11.82, and dmft in children aged 3-6 years was 7.7. These results show a slight improvement in oral hygiene, restoration, and prevention in urban than in rural parts of Croatia. In a study that analyzed oral health in the capital cities of the former Yugoslavia in 1986, mean dmft in children aged 6 years was 8.4 and mean DMFT 0.9 (34). Mean DMFT in 6-year-old children was 0.8 and mean DMFT in 12-year-old children was 5.7 (34). The study conducted in Zagreb in 1992 reported a mean DMFT of 2.48 and 3.75 in children aged 7-8 and 9-10 years, respectively (35). Although these studies are 20-25 years old, our results showed only limited improvement in oral indexes, which is unsatisfactory for such a long time period and indicates that all new preventive and restorative measures have had limited effectiveness.

This study showed the importance of epidemiological studies for caries prevention and maintenance of oral health. The percentage of children with one or more caries lesions was too high considering the currently available preventive possibilities. The obtained DMFT and DMFS indexes, in comparison with other countries, illustrate poor oral health, low oral hygiene, and unsatisfactory prevention in Croatia, as well as a need to invest in modern preventive and therapeutic methods.

Funding This study is supported by the City Office for Health and War Veterans of Zagreb, Croatia. It was performed within the research project "Prevention of Early Caries in Children: Clinical Evaluation of Preventive Methods" No. 065-0653147-2056 approved by the Ministry of Science and Technology of the Republic of Croatia.

Ethical approval received from the Ethics Committee of the Zagreb School of Dental Medicine.

Declaration of authorship WD was involved in formulation of the basic concept of this study, statistical analysis and writing of the manuscript; he substantially contributed to study conception and design as well as data acquisition, analysis, interpretation, critical revision, and final approval of the manuscript. BD was involved in clinical examinations of participants, data collection, arranging the tables, revising the manuscript, including translation and language style. OLD was involved in collection of the participants, designing the study, and managing the prevention project and manuscript preparation.

Competing interests All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organization for the submitted work; no financial relationships

with any organizations that might have an interest in the submitted work in the previous 3 years; no other relationships or activities that could appear to have influenced the submitted work.

References

- 1 Ismail AI. Prevention of early childhood caries. *Community Dent Oral Epidemiol.* 1998;26:49-61. [Medline:9671200](#)
- 2 Davies GN. Early childhood caries – a synopsis. *Community Dent Oral Epidemiol.* 1998;26:106-16. [Medline:9671208](#)
- 3 Rosenblatt A, Zarzar P. The prevalence of early childhood caries in 12- to 36-month-old children in Recife, Brazil. *ASDC J Dent Child.* 2002;69:319-24, 236. [Medline:12613320](#).
- 4 Filstrup SL, Briskie D, da Fonseca M, Lawrence L, Wandera A, Inglehart MR. Early childhood caries and quality of life: child and parent perspectives. *Pediatr Dent.* 2003;25:431-40. [Medline:14649606](#)
- 5 Sheiham A. Oral health, general health and quality of life. *Bull World Health Organ.* 2005;83:644. [Medline:16211151](#)
- 6 Iijima Y, Takagi O. In situ acid resistance of in vivo formed white spot lesions. *Caries Res.* 2000;34:388-94. [Medline:11014905](#) [doi:10.1159/000016613](#)
- 7 Hicks MJ, Flaitz CM. Epidemiology of dental caries in the pediatric and adolescent population: a review of past and current trends. *J Clin Pediatr Dent.* 1993;18:43-9. [Medline:8110613](#)
- 8 Chow LC, Vogel GL. Enhancing remineralization. *Oper Dent.* 2002;26 suppl 6:27-38.
- 9 Wyne AH. Caries prevalence, severity, and pattern in preschool children. *J Contemp Dent Pract.* 2008;9:24-31. [Medline:18335116](#)
- 10 National Caries Program-NIDR. The prevalence of dental caries in United States children, 1979-1980. NIH Publication No. 82-2245;1981.
- 11 Pizzo G, Piscopo MR, Matranga D, Luparello M, Pizzo I, Giulia G. Prevalence and socio-behavioral determinants of dental caries in Sicilian schoolchildren. *Med Sci Monit.* 2010;16:PH83-9. [Medline:20885361](#)
- 12 Campus G, Lumbau A, Lai S, Solinas G, Castiglia P. Socio-economic and behavioural factors related to caries in twelve-year-old Sardinian children. *Caries Res.* 2001;35:427-34. [Medline:11799283](#) [doi:10.1159/000047486](#)
- 13 Vadiakas G, Oulis CJ, Tsinidou K, Mamai-Homata E, Polychronopoulou A. Socio-behavioural factors influencing oral health of 12 and 15 year old Greek adolescents. A national pathfinder survey. *Eur Arch Paediatr Dent.* 2011;12:139-45. [Medline:21640058](#)
- 14 Gatou T, Koletis Kounari H, Mamai-Homata E. Dental caries prevalence and treatment needs of 5- to 12-year-old children in relation to area-based income and immigrant background in Greece. *Int Dent J.* 2011;61:144-51. [Medline:21692785](#) [doi:10.1111/j.1875-595X.2011.00031.x](#)
- 15 Muratbegovic AA, Markovic N, Zukanovic A, Kobaslija S, Dragas MS, Juric H. Oral health related to demographic features in Bosnian children aged six. *Coll Antropol.* 2010;34:1027-33. [Medline:20977098](#)
- 16 Zukanovic A, Muratbegovic A, Kobaslija S, Markovic N, Ganibegovic M, Beslagic E. Relationships between socioeconomic backgrounds, caries associated microflora and caries experience in 12-year-olds in Bosnia and Herzegovina in 2004. *Eur J Paediatr Dent.* 2008;9:118-24. [Medline:18844440](#)
- 17 Djurickovic M, Ivanovic M. The state of oral health in children at the age of 12 in Montenegro [in Serbian]. *Vojnosanit Pregl.* 2011;68:550-5. [Medline:21899174](#) [doi:10.2298/VSP1107550D](#)
- 18 Begzati A, Meqa K, Siegenthaler D, Berisha M, Mautsch W. Dental health evaluation of children in Kosovo. *Eur J Dent.* 2011;5:32-9. [Medline:21228954](#)
- 19 Bravo Pérez M, Casals Peidró E, Cortés Martinicorena FJ. Oral health survey of Spain 2005 [in Spanish]. *RCOE.* 2006;11:409-56.
- 20 Bravo M, Cortés J, Casals E, Llena C, Almerich-Silla JM, Cuenca E. Basic oral health goals for Spain 2015/2020. *Int Dent J.* 2009;59:78-82. [Medline:19489286](#)
- 21 Borutta A, Brauner K, Hufnagl S, Marton S, Mavrodisk Z, Tarjan I. Oral health in 8-9 year-old children in Saxony-Anhalt (Germany) and in two Hungarian cities (Budapest and Debrecen). *Community Dent Health.* 2006;23:26-30. [Medline:16555716](#)
- 22 Croatian Bureau of Statistics. Census 2011. Available from: http://www.dzs.hr/Hrv_Eng/publication/2011/SI-1441.pdf. Accessed: October 15, 2011.
- 23 City Bureau for Urban Planning and Environmental Protection – Statistics Department. Population, households and dwellings 2001 [in Croatian]. Available from: http://www1.zagreb.hr/zgstat/documents/gradske_cetvrti_gz_stanovnistvo_kucan_stvstanovi_2001/publikacija.pdf. Accessed: October 15, 2011.
- 24 City Bureau for Urban Planning and Environmental Protection – Statistics Department. Population of Zagreb according to sex and age, 2001 [in Croatian]. Available from: http://www1.zagreb.hr/zgstat/documents/stanovnistvo_gz_%20prema_dobnoj_spolno_strukturi_2001/publikacija.pdf. Accessed: October 15, 2011.
- 25 World Health Organization. Oral Health Surveys, basic methods. 4. edition. Geneva (Switzerland): World Health Organization; 1997.
- 26 World Health Organization. The World Oral Health Report, 2003. Continuous improvement of oral health in the 21 st century – the approach of the WHO Global Oral Health Programme. Geneva (Switzerland): World Health Organization; 2003.
- 27 Hobdell MH, Myburgh NG, Kelman M, Hausen H. Setting global goals for oral health for the year 2010. *Int Dent J.* 2000;50:245-9. [Medline:15988881](#) [doi:10.1111/j.1875-595X.2000.tb00560.x](#)
- 28 Marthaler TM, O'Mullane DM, Vrbic V. The prevalence of dental caries in Europe 1990-1995. ORCA Saturday afternoon symposium 1995. *Caries Res.* 1996;30:237-55. [Medline:8773416](#) [doi:10.1159/000262332](#)
- 29 Marthaler TM. Changes in dental caries 1953-2003. *Caries Res.*

- 2004;38:173-81. [Medline:15153686](#) [doi:10.1159/000077752](#)
- 30 Almerich-Silla JM, Montiel-Co JM. Influence of immigration and other factors on caries in 12- and 15-yr-old children. *Eur J Oral Sci.* 2007;115:378-83. [Medline:17850426](#) [doi:10.1111/j.1600-0722.2007.00471.x](#)
- 31 Bratthall D. Introducing the Significant Caries Index together with a proposal for a new global oral health goal for 12-year-olds. *Int Dent J.* 2000;50:378-84. [Medline:11197197](#) [doi:10.1111/j.1875-595X.2000.tb00572.x](#)
- 32 Campus G, Sacco G, Cagetti M, Abati S. Changing trend of caries from 1989 to 2004 among 12-year old Sardinian children. *BMC Public Health.* 2007;7:28. [Medline:17331258](#) [doi:10.1186/1471-2458-7-28](#)
- 33 Juric H, Klaric T, Zagar M, Bukovic D Jr, Jankovic B, Spalj S. Incidence of caries in children of rural and subrural areas in Croatia. *Coll Antropol.* 2008;32:131-6. [Medline:18494198](#)
- 34 Vrbic V, Vulovic M, Rajic Z, Topic B, Tatic E, Malic M, et al. Oral health in SFR Yugoslavia in 1986. *Community Dent Oral Epidemiol.* 1988;16:286-8. [Medline:3180717](#) [doi:10.1111/j.1600-0528.1988.tb01776.x](#)
- 35 Rošin-Grget K, Lulić Dukić O, Radionov D, Jelušić-Franulović B, Linčir I. Dental health of children in primary school "Retkovec" in Zagreb [in Croatian]. *Acta Stomatol Croat.* 1992;26:61-6.