

Supplementary Table 1. Differences in clinical characteristics and medications between male and female gender in ACS and sCAD groups

	ACS			sCAD		
	Male n=129	Female n=58	<i>P</i>	Male n=46	Female n=26	<i>P</i>
Age (years); median (IQR 25-75 th) ¹	62.0 (54.0-72.0)	75.0 (67.0-79.0)	<0.001	62.5 (55.0-70.0)	67.5 (59.0-75.0)	0.029
Diabetes mellitus; n (%) ²	17 (13.2)	10 (17.2)	0.465	11 (23.9)	11 (42.3)	0.104
Hyperlipoproteinaemia; n (%) ²	36 (27.9)	14 (24.1)	0.590	21 (45.7)	12 (46.2)	0.967
Smoking cigarettes; n (%) ²	57 (44.2)	11 (19.0)	0.001	9 (19.6)	7 (26.9)	0.471
MAP (mmHg); median (IQR 25-75 th) ¹	90.0 (81.7-96.7)	86.7 (78.3-96.7)	0.260	96.7 (90.0-103.3)	100.0 (95.0-110.0)	0.046
BMI (kg/m ²); median (IQR 25-75 th) ¹	27.7 (25.1-30.3)	29.3 (25.7-32.9)	0.098	28.5 (25.1-29.9)	25.8 (23.3-30.8)	0.305
LV EF (%); median (IQR 25-75 th) ¹	50.0 (48.0-55.0)	50.0 (45.0-55.0)	0.414	58.0 (55.0-60.0)	60.0 (50.0-60.0)	0.865
Number of segment; median (IQR 25-75 th) ¹	2.0 (1.0-3.0)	3.0 (2.0-4.0)	0.079	2.0 (2.0-4.0)	2.0 (1.0-3.0)	0.755
Gensini score; median (IQR 25-75 th) ¹	47.0 (32.0-80.0)	50.5 (34.0-80.0)	0.376	25.0 (11.0-37.0)	30.0 (9.0-49.0)	0.783
ASA before hospitalization; n (%) ²	18 (14.0)	22 (37.0)	<0.001	29 (63.0)	19 (73.1)	0.386
Statins before hospitalization; n (%) ²	15 (11.6)	9 (15.5)	0.500	25 (54.3)	10 (38.5)	0.062

¹ Mann-Whitney U test, ² Chi-square test

IQR 25-75th, interquartile range 25-75th percentiles; MAP, mean arterial pressure; BMI, body mass index; LV EF, left ventricular ejection fraction; ACS, acute coronary syndrome; sCAD, stable coronary artery disease; ASA, acetyl salicylic acid