



Reference Resources

Caveats from AAALAC's Council on Accreditation regarding this resource:

Report of the ACLAM Task Force on Rodent Euthanasia*.

Artwohl J, Brown P, Corning B, Stein S. ACLAM Task Force.

JAALAS. 45(1):98-105, 2006.

*This reference was adopted by the Council on Accreditation with the following caveats:

Hypothermia is identified as an accepted method for inducing anesthesia in rodent fetuses (B. Euthanasia of Fetuses 2.d.) or pups 6 days of age or less (C. Euthanasia of Neonates 1.f.). This practice is common in the United States, but may not be accepted internationally. Suitability of this technique should be determined on a case-by-case basis with consideration of input from the Attending Veterinarian and the IACUC or other oversight body.

The science surrounding the use of carbon dioxide for rodent euthanasia is evolving. At present, there is no universally accepted procedure for euthanizing rodents using carbon dioxide. Participating institutions should be aware that updates to the science may result in refined methodology and may alter the Council on Accreditation's view of the methods described in this report.

This Reference Resource begins on the next page....

Public Statement

Report of the ACLAM Task Force on Rodent Euthanasia

James Artwohl, Patricia Brown, Brian Corning and Susan Stein

The ACLAM Task Force on Rodent Euthanasia was appointed by President Lynn Anderson in 2002 in response to growing concerns and controversy regarding techniques that were commonly used for rodent euthanasia. Three issues were targeted as the focus of the report: euthanasia of fetal and neonatal rodents, the use of carbon dioxide for rodent euthanasia, and the impact of euthanasia techniques on data. The charge to the Task Force was to create a document that summarized in a scholarly and comprehensive manner all available data-based literature relevant to these topics, to assess the scientific merit of the design and conclusions of those studies, and to compile valid information into a concise and cohesive document that could serve as a resource for diplomates, other veterinarians, IACUC members, regulatory bodies, and research scientists.

The Task Force has fulfilled this charge in an exemplary manner. During 2004-2005, the ACLAM officers and Board of Directors (BOD) reviewed and critiqued 2 draft versions of the report, and suggestions for change were incorporated into the document presented here. In July 2005, the BOD voted to forego the usual process of distributing the document to the ACLAM membership for comment before release based on 2 considerations. First, the literature relevant to rodent euthanasia is continually expanding. As such, at each revision, the Task Force was compelled to incorporate new data and citations. Their consensus view was that new data would continue to emerge, and the document would require continual revision as the review process continued. Related to that, the 2nd consideration of the BOD was that information already accumulated would be of immediate utility to the stake-holders listed above.

In lieu of a pre-publication comment period, the BOD and the Task Force instead invite all diplomates, as well as other parties, to comment via email or mail to the BOD liaison for this project, who will compile and maintain all remarks. After an interval deemed appropriate by the ACLAM President, a 2nd Task Force will be appointed to update and modify the Report. Comments will be considered at that time.

I want to personally thank all members of the Task Force for their conscientious and comprehensive efforts in compiling this information. They have created a valuable and informative synthesis that should serve as a resource to the community for years to come.

—Linda A Toth, DVM, PhD
ACLAM BOD Liaison to Task Force on Rodent Euthanasia

Introduction

The guidelines below were prepared by the American College of Laboratory Animal Medicine (ACLAM) to expand upon the information provided by the Report of the AVMA Panel on Euthanasia with regard to euthanasia of rodents in biomedical settings. Database searches were designed with assistance from a library scientist with advanced degrees in public health, education and research, training by AWIC and responsibility for veterinary reference materials. Keywords were selected to include all ages of rodents and all categories of rodent euthanasia identified by the 2000 AVMA Panel on Euthanasia.² Peer reviewed publications from 1912 to 2005 were identified and evaluated prior to inclusion as references.

Professional consultation with the attending veterinarian is essential when developing plans for euthanasia and when applying these guidelines. The intent of this document is to provide guidance on rodent euthanasia performed at biomedical research facilities. Selection of the optimal euthanasia methods must be assessed on an individual basis; however, there should be consistency in the goals of all methods employed. The euthanasia of rodents should be humane, minimizing pain and fear, delivered in accordance with current regulations, ensure rapid onset of unconsciousness followed by

death, and avoid risk and aversion for animals and personnel. Specific information on fetal and neonatal euthanasia, use of CO₂ as an euthanasia agent, and the influence of euthanasia methods on frequently measured scientific parameters should be incorporated into the institution's educational program for investigators.

Euthanasia of Fetal and Neonatal Rodents

A. Background

The Report of the AVMA Panel on Euthanasia² provides limited recommendations for the euthanasia of prenatal or neonatal animals and no specific recommendations on altricial or precocial rodents. The report states: "When ovarian hysterectomies are performed, euthanasia of feti should be accomplished as soon as possible after removal from the dam." It also states "Neonatal animals appear to be resistant to hypoxia, and because all inhalant agents ultimately cause hypoxia, neonatal animals take longer to die than adults."² The Panel recommends "inhalant agents not be used alone in animals less than 16 weeks old except to induce loss of consciousness, followed by the use of some other method to kill the animal."²

The current scientific literature provides limited evidence for the effectiveness of any of the recommended rodent euthanasia

methods when performed on fetuses or neonates or the outcome on the fetuses when performed on the pregnant mother.

B. Euthanasia of Fetuses

By the 3rd trimester of gestation, the neural tube has developed into a functional brain, and the likelihood that a fetus may perceive pain should be considered.^{12,30} No definitive evidence indicates that prenatal rodents perceive pain, but reflexive behavior observed in fetal animals correlates with adult responses to painful stimuli.^{15,55} However, low arterial oxygen concentrations may limit higher cortical processing that would mediate fetal arousal and awareness.⁴³

1. Mouse, Rat, and Hamster Fetuses up to 15 Days' and Guinea Pig Fetuses up to 35 Days' Gestation
 - a. Neural development during this developmental stage is minimal and pain perception is considered unlikely.^{36,73}
 - b. Euthanasia of the mother or removal of the fetus should ensure rapid death of the fetus due to loss of blood supply and non-viability of fetuses at this stage of development.³⁷
2. Mouse, Rat, and Hamster Fetuses over 15 Days' and Guinea Pig Fetuses over 35 Days' Gestation through Birth
 - a. The neural development during this developmental period supports the likelihood that pain may be perceived.^{30,36,73} Observations of near-term mouse and rat fetuses in vivo indicate behavioral responses to sensory stimulation.^{17,65}
 - b. Methods of euthanasia of fetuses
 - i. Skillful injection of chemical anesthetics in sufficient quantities to ensure death.
 - ii. Decapitation with sharp surgical scissors or cervical dislocation.
 - c. Rapid freezing in liquid nitrogen without prior anesthesia is not considered to be humane.²
 - d. When chemical fixation of the whole fetus is required, fetuses should be anesthetized prior to immersion in, or perfusion with, fixative solutions. Anesthesia may be induced by hypothermia,^{19,54} or by injection with a chemical anesthetic.⁶⁷
 - e. Rodent fetuses are resistant to hypoxia.⁶³ Near-term rat fetuses experiencing umbilical cord occlusion exhibited respiratory movements for up to 40 min after occlusion.⁵⁸ Fetuses require extended exposure to inhalant anesthetics, including CO₂.³⁷
 - f. When fetuses are not required for study, the method chosen for euthanasia of a pregnant mother should ensure cerebral anoxia to the fetus and minimally disturb the uterine milieu to minimize fetal arousal.^{37,43} A recommended method for euthanasia of the mother is CO₂ exposure followed by cervical dislocation.⁴⁷

C. Euthanasia of Neonates

1. Mouse, Rat, and Hamster Neonates up to 10 Days of Age
 - a. Maturation of nociceptors and the development of excitatory and inhibitory receptor systems occur during the period just prior to birth and extend into the 2 wk of postnatal life.^{26,28,57,70}
 - b. Resistance to hypoxia results in a prolonged time to unconsciousness when CO₂ inhalation is used as a euthanasia agent.^{2,37,47} The duration of exposure to

carbon dioxide varies with the age of the neonate. Inbred and outbred neonatal mice less than 7 d of age may differ in susceptibility to CO₂, requiring exposures as long as 50 min to ensure euthanasia.⁵⁵

- c. When using CO₂ for euthanasia, death must be verified prior to disposal of the carcass.⁵¹
 - d. Other methods for the euthanasia of neonatal mice and rats
 - i. Injection of chemical anesthetics in sufficient quantities to ensure death.
 - ii. Decapitation.
 - iii. Cervical dislocation.
 - e. Immersion in liquid nitrogen should be performed only if preceded by anesthesia. Anesthesia should precede immersion in, or perfusion with, chemical fixatives.
 - f. Anesthesia in neonatal rodents may be induced by inhalant or injectable anesthetics. Prolonged exposure to inhalant anesthetics (e.g., halothane or isoflurane) may be necessary. Alternatively, hypothermia may be used to induce anesthesia in pups 6 d of age or less.^{19,54} The attending veterinarian should be consulted for appropriate techniques and drug dosages.
2. Guinea Pig Neonates
Follow guidelines for adults.²
 3. Mouse, Rat, and Hamster Neonates over 10 Days of Age
Follow guidelines for adults.²

The Use of CO₂ for Euthanasia of Rodents

A. Background

Carbon dioxide (CO₂) is a frequently used euthanasia agent for small laboratory animals due to its rapid onset of action, safety, low cost, and ready availability. It most commonly is used to euthanize rats and mice, which constitute the majority of animals used in biomedical research and are the focus of most studies on the use of CO₂ euthanasia. The same delivery system and equipment can be used to euthanize either single animals or groups of animals with CO₂. Despite its widespread use, euthanasia methods using CO₂ are not standardized. The current peer reviewed literature does not establish consistent requirements for CO₂ euthanasia and or even provide a clear definition of what constitutes a humane death. The acceptability of CO₂ for euthanasia under various conditions, and for various species and ages of animals, must continue to be re-evaluated as new data become available.

B. General Considerations

Changes in the animal's environment or novel conditions should be minimized to the degree that is practical. Rodents are sensitive to their environment and to handling.^{42,59,62} Removal from the home cage,¹⁴ regrouping with other animals,⁴⁴ introduction to new sites and odors,¹¹ and transport and placement into the euthanasia chamber can alter physiologic and metabolic parameters and possibly cause stress. Researchers and animal care staff should seek methods that minimize the stress experienced by rodents that undergo CO₂ euthanasia. Transporting animals and performing euthanasia in the home cages, using carts that are quiet, roll freely, and do not jostle cages or occupants, and minimizing regrouping to prevent social aggression are simple approaches to lessening potentially stressful conditions.

C. Euthanasia Chambers

1. Euthanasia chambers should be kept clean and free of debris and excreta.
2. The euthanasia chamber should be large enough to permit each animal to stand on the floor of the chamber with all 4 feet and have sufficient space to turn around and perform normal postural adjustments.

D. CO₂ Gas Delivery Systems

1. Sufficient carbon dioxide must be introduced into the chamber to totally displace the residual air by both mixing and dilution. Ideally, the inlet for delivery of CO₂ and any diffusion devices in the euthanasia chamber should provide a predictable and controllable elevation in CO₂ concentration.
2. Excess gas must be allowed to escape from the chamber in a way that allows a gradual increase in the concentration of CO₂ at the floor of the container that holds the animal. Escape of the gas mixture through a port, or other opening at the top of the chamber, must occur in a controlled manner that neither pressurizes the chamber nor permits reflux of room air into the chamber.
3. Carbon dioxide should be delivered using a 2-stage regulator, with the 2nd stage capable of adjustable fixed flow rates.
4. Large chambers designed for euthanasia of groups of animals may require multiple inlets, or diffusion devices, to facilitate different configurations for CO₂ introduction.
5. The use of heated valves assures constant delivery of gas to the chamber by avoiding the formation of dry ice within valves and regulating systems when units are used for prolonged or repeated periods.
6. The filling rate of the chamber should be based upon the time required to rapidly and successfully render the animals unconscious. This may differ from the amount of time required to achieve a lethal concentration of CO₂. Each type of chamber will have a different CO₂ filling profile based upon gas flow rate, gas dispersion characteristics, gas inlet locations and chamber dimensions.
7. Chambers should be filled with CO₂ at a flow rate that balances the time to unconsciousness with such associated aversive stimuli as noise or high velocity air movement.
8. A fill rate of 20% of the chamber volume per minute has been recommended as an appropriate means to achieve a lethal concentration.³¹ However, animals should be closely observed during the filling process, as individual systems may require adjustment to achieve the desired effect.²

E. Pre-filling vs Not Pre-filling the Euthanasia Chamber

1. Because inspiration of high concentrations of CO₂ is both aversive and painful,^{9,19,40} a recommended procedure is to place animals into a chamber that contains room air and then to gradually introduce CO₂.
2. The use of CO₂/oxygen gas mixtures and slow fill rates prolong the time to unconsciousness and death and may

increase distress for the animals. There is no conclusive evidence that adding pure oxygen to carbon dioxide makes this procedure less stressful to animals.^{13,20,29,39} A fill rate of 20% of the chamber volume per minute with carbon dioxide, added to existing room air in the chamber should be appropriate to achieve a balanced gas mixture to fulfill the objective of rapid unconsciousness with minimal distress to the animals.

F. Cautionary Information

1. Animal carcasses should not be exposed to room air until death has ensued with high certainty, as the anesthetic effects of CO₂ can be quickly reversed in the presence of oxygen.
2. Individual rodents may become apneic at certain concentrations of CO₂, giving the false impression that death has occurred.⁹
3. Confirmation of death should be based not on a single sign, such as cessation of breathing, but on multiple signs, such as physical examination, exposure to room air (under observation), or adjunctive methods of euthanasia (decapitation, cervical dislocation, pneumothorax).
4. Euthanasia apparatus should be regularly evaluated to ensure proper functionality sufficient to achieve 100% euthanasia of all animals. Failure to function correctly may result in the need to re-expose an animal to carbon dioxide to achieve euthanasia. Re-exposure should take place before the animal regains consciousness.

The Influence of Euthanasia Method on Frequently Measured Scientific Parameters

A. Background

The method of euthanasia can influence the validity of scientific results. *The Guide for the Care and Use of Laboratory Animals* indicates the appropriate euthanasia method depends on many criteria, including compatibility with research objectives.⁴⁸ It further states, "The selection of specific agents and methods for euthanasia will depend on the species involved and the objectives of the protocol."⁴⁸ Euthanasia, as a process, separates the presentation of new variables, treatments or environmental changes to the living system from the terminal collection of tissues and blood for additional study or analysis. In itself, the euthanasia method can alter physiologic parameters and responses.

The effects of handling during the euthanasia process, proficiency of personnel performing euthanasia, and mechanical efficiency of equipment can introduce variables that influence the welfare of the animal and the interpretation of the data. Such factors as the species and age of animal, measurements to be assessed, sampling sites, and time of tissue collection, additionally influence sample analysis and histology.

The existing literature should be assessed for general information. However, the specific impact of any euthanasia method on scientific results may require case-by-case validation.

The euthanasia technique should minimally impact the welfare of the animal and the handler and must support collection of reproducible scientific data.

The researcher must evaluate the scientific consequences of the chosen method of euthanasia.

B. Biological Effects of Euthanasia Techniques

Table 1. Biologic effects of decapitation^{3,5,16,49,56,60,66}

Effect	Mechanism
Increase in plasma sodium	Hemolysis
Increase in plasma potassium	
Increase in GABA concentrations (brain)	
Increase in Alanine (brain)	
Increase in plasma ascorbic acid (30-40% > resting state)	
Increase in blood catecholamine levels	Continued postmortem neurochemical alterations
Increased plasma calcium, magnesium	Stress stimulus → mobilization from tissues to blood; generalized metabolic response secondary to sympathoadrenal response some handling related stimulation.
No change in vasoactive intestinal peptides (brain)	
No change in neuropeptide Y (brain)	
Alteration in rat heart mitochondria function	
Increase in serum corticosterone	
	Possible handling stress

Table 2. Effects of physical and pharmacological euthanasia methods

Method	Physiologic effect
Methoxyflurane and decapitation ¹⁰	Increase in prostacyclin (vasodilator that inhibits platelet aggregation) Vascular contractility suppressed Decreased vascular contractility
Ether and decapitation, or decapitation alone ⁵⁰	No statistical difference in prolactin levels or LH/FSH secretory properties of cultured anterior pituitary cells
Ether and decapitation ⁷⁴	No change in estrogen receptors/progesterone receptors in rat uteri
Ketamine and decapitation ^{50,74}	No change in estrogen receptors/progesterone receptors in rat uteri
Pentobarbital and decapitation ⁴	Increase in acetylcholine release in the brain
Halothane and decapitation ²¹	Increase in plasma ascorbic acid Increase in plasma catecholamines

Table 3. Effects on reproductive hormones: The following combinations may be unsuitable for studies of serum androgens

Decapitation in combination with agents listed below ^{49,71}	Male rats								Mechanism: direct effect on testes	
	Immature				Mature				Circulating Androstenedione	
	LH	FSH	Prolactin	Testosterone	LH	FSH	Prolactin	Testosterone	Castrated	Intact
Xylazine	-	-		↓	-	-	↑	↓		↓ or -
Biotol	-	-			-	-		↓		↓ or -
Thiopental	-	-			-	-		↓		↓ or -
Pentobarbital	-	-		↓	-	-	↑	↓		↓ or -
Ketamine	↓	↓		↓	-	-		↓	↑	↓ or -
Halothane	↓	↓		↓	-	-		↓		↓ or -
Ether (tested on castrated rats)	↑	↑	↑	↓	-	-		↓		↓ or -

↓ = decreased ↑ = increased - = no change.

Table 4. Biologic effects of euthanasia induced by pharmacologic and/or physical methods

Method of euthanasia	Effect	Mechanism
Injectable Pentobarbital ^{5,53,61} a,b	Decreased muscular contractility in isolated muscle preps Decreased GI smooth muscle contractility when given orally or intravenously; not seen in intraperitoneal route Intraperitoneal administration causes increased colonic contractility in response to acetylcholine Decreased spontaneous and drug induced vascular smooth muscle contractility Decreased catecholamine levels Increased partial pressure of CO ₂ in arterial blood Increased serum activity renin Increased plasma aldosterone Splenic enlargement Increased plasma glucose and insulin Increased liver glycogen Decreased plasma triglycerides Increase in plasma insulin	Decreased calcium transport Increased CO ₂ in arterial blood may change blood pH, which then changes metabolic indices Increased glucose production or decreased glucose clearance
Cervical dislocation/ cervical fracture ^{32, 68, 72}	Decreased coronary flow; decreased contractile function in isolated perfused heart preparations Normal lymphocyte proliferation High levels of serotonin in lung Increase in granulocyte and macrophage colony forming cell counts in murine bone marrow cultures	Possible decreased sensitivity of B-adrenergic receptors secondary to cervical fracture Entrapment of platelets in pulmonary capillaries Apparent alteration of marrow stem cell pool
Cervical dislocation and methoxyflurane ³²	Increased mitogen induced lymphocyte proliferation Normal cytolytic T lymphocytes (CTL) response	
Cervical dislocation and pentobarbital ³²	Increased mitogen induced lymphocyte proliferation Decreased CTL response	
Cervical dislocation and halothane ³²	Normal mitogen induced lymphocyte proliferation Decreased CTL response	
CO ₂ and cervical dislocation ³²	Normal mitogen induced lymphocyte proliferation Decreased CTL response	
CO ₂ and decapitation ^{4,23,66}	Normal LH, FSH, prolactin, corticosterone Activity of cholinergic markers identical to decapitation only Altered GABA _A receptor function	
Focused beam microwave irradiation (FBMI) ^{41,45} c	Best technique for measuring adenosine levels Decreased brain amino acids: alanine, GABA, ethanolamine, NH ₃ , valine, leucine, isoleucine, tyrosine, phenylalanine, glycine, aspartate Increased levels of reduced glutathione, glutamate 5 fold decrease in d prostaglandin and Thromboxane B ₂ (mouse brain) Twice the concentration of substance P, neurokinin A, and neurotensin in brain tissue compared to decapitation	Due to rapid inactivation of metabolizing enzyme Possible enzyme inactivation by microwave irradiation causing increased recovery of peptides Possible disintegration of neuropeptide containing tissue compartments, or decreased binding of carrier proteins, releasing more peptides
CO ₂ ^{8,52,69}	100% CO ₂ : decreased mean corpuscular hemoglobin (NP) ^d Increased total leukocytes and granulocytes (P) ^e Decreased liver glycogen, pyruvate, ATP No change in platelet counts	CO ₂ causes acidosis that affects RBC parameters
CO ₂ or CO ₂ /O ₂ ^{8,27,34,46,52,69}	Increased hematocrit, mean corpuscular volume No change in serum norepinephrine, dopamine, serotonin, corticosterone ^f Decreased serum creatine kinase, aspartate aminotransferase Significant decreased liver glycogen stores Increased serum glucose Decreased activity of enzymes regulating branched chain amino acid degradation Decreased mean erythrocyte hemoglobin, mean corpuscular hemoglobin concentration	 CO ₂ causes acidosis that produces stimulation of enzymes of the glycolytic pathway
70% CO ₂ /30% O ₂ vs 100% (Pre-charged) ⁵²	Decreased number of circulating CD3 ⁺ and CD8 ⁺ T cells Increase in CD10 ⁺ B cells in circulation	
70% CO ₂ /30% O ₂ vs 100% (Not pre-charged) ⁵²	Increased number of circulating CD3 ⁺ , CD4 ⁺ , and CD8 ⁺ T cells	NOTE: 100% CO ₂ (Non-precharged and pre-charged) had overall greater T-cell counts than 70% CO ₂ /30% O ₂ euthanized animals
Isoflurane ⁸	No change in liver glycogen	

^apreferred for isolated beating heart preparations.

^bpreferred method (by IV route) for collection of tissues, including liver, for cyclic AMP assay.

^cFBMI minimizes post-mortem neurochemical changes. Leaves brain landmarks intact.

^d(NP) = Not precharged.

^e(P) = Precharged.

^fexposure to CO₂ prior to decapitation.

Table 5. Anesthetics – ketamine hydrochloride, pentobarbital, chloral hydrate, chloralose and halothane in combination

Fructose -2-6-biphosphate ³⁵	Significant increase in brain, heart, skeletal muscle concentrations
---	--

reported to release massive sympathetic response with ↑ catecholamines from adrenal gland.

Table 6. Gross/histopathology changes^{1,24,25,33,64}

Ether	Decapitation	CO ₂ ^a	Methoxyflurane	Pentobarbital	Physical Methods (DC, CD)	Methods Listed in this Chart
Lung: interstitial edema, marked alveolar emphysema	Lung: emphysema, hemorrhage, blood in alveolar spaces	Lung: congestion, hemorrhage, emphysema, atelectasis; Cardiac muscle: variable degenerative changes (influenced by time of exposure to CO ₂ causing acidosis, hypoxia) CO ₂ + O ₂ Lung: severe edema and hemorrhage, extravasation to alveoli Cardiac muscle: variable degenerative changes (influenced by time of exposure to CO ₂ causing acidosis, hypoxia), capillary bleeding causing marked extravasation of blood	Lung: congestion Spleen: splenomegaly	Lung: emphysema congestion Spleen: emphysema, congestion GI serosa: emphysema, congestion Cardiac muscle: Acute degenerative lesions Kidney cortex: circulatory changes Other: Peritoneal congestion, sanguinous fluid in abdominal cavity	Lung: emphysema, bleeding Neck/Brain: local tissue trauma	No change in sperm motion

NOTE: DC (decapitation), CD (cervical dislocation), CO₂, Intracardiac pentobarbital more suitable for histology of abdominal viscera.

^aproduces changes in hemodynamics—capillary contraction, followed by dilation of capillaries and veins (except lung vessels); depresses cerebral cortex, stimulates chemoreceptors; extravasation to alveoli: Not seen in all rodent species.

Table 7. Additional Factors that Influence the Outcome of Euthanasia^{6,7,18,22,38,56}

1. Handling: May cause sympathoadrenal discharge, which affects plasma glucose, progesterone plasma catecholamines. Habituating the animals to handling may mitigate this effect.
2. Environmental stimuli (for example, noise) can increase plasma corticosterone concentrations.
3. Sequence: The order of euthanasia for rats housed in pairs produced significant differences in plasma tryptophan and unesterified fatty acids, plasma corticosterone, plasma protein lactate levels, substance P, cholecystokinin, somatostatin.

References

1. **Ambrose N, Wadham J, Morton D.** 2000. Refinement of euthanasia. Progress in the reduction, refinement and replacement of animal experimentation. Elsevier Science. p 1159–1170.
2. **Beaver BV, Reed W, Leary S, McKiernan B, Bain F, Schultz R, Bennett BT, Pascoe P, Shull E, Cork LC, et al.** 2001. Report of the AVMA panel on euthanasia. *J Am Vet Med Assoc* **218**:688.
3. **Behrens WA, Madere R.** 1979. Effects of handling, anesthesia and decapitation on plasma ascorbic acid in the rat. *Nutr Rep Int* **19**:419–426.
4. **Berger-Sweeney J, Berger UV, Sharma M, Paul CA.** 1994. Effects of carbon dioxide-induced anesthesia on cholinergic parameters in rat brain. *Lab Anim Sci* **44**:369–371.
5. **Bhathena SJ.** 1992. Comparison of effects of decapitation and anesthesia on metabolic and hormonal parameters in Sprague-Dawley rats. *Life Sci* **50**:1649–1655.
6. **Bickhardt K, Buttner D, Muschen U, Plonait H.** 1983. Influence of bleeding procedure and some environmental conditions on stress-dependent blood constituents of laboratory rats. *Lab Anim* **17**:161–165.
7. **Brodin E, Rosen A, Schott E, Brodin K.** 1994. Effects of sequential removal of rats from a group cage and of individual housing of rats on substance P, cholecystokinin and somatostatin levels in the periaqueductal grey and limbic regions. *Neuropeptides* **26**:253–60.
8. **Brooks SPJ, Lampi BJ, Bijun CG.** 1999. The influence of euthanasia methods on rat liver metabolism. *Contemp Top Lab Anim Sci* **38**(6):19–24.
9. **Britt DP.** The humaneness of carbon dioxide as an agent of euthanasia for laboratory rodents. Euthanasia of Unwanted, Injured or Diseased Animals for Educational or Scientific Purposes. Hertfordshire (UK): Universities Federation for Animal Welfare.
10. **Butler MM, Griffey SM, Clubb FJ, Gerrity L, Campbell WB.** 1990. The effect of euthanasia technique on vascular arachidonic acid metabolism and vascular and intestinal smooth muscle contractility. *Lab Anim Sci* **40**:277–83.
11. **Carr VM, Menco B, Yankova MP, Morimoto RI, Farbman AI.** 2001. Odorants as cell-type specific activator of a heat shock response in the rat olfactory mucosa. *J Comp Neurol* **432**:425–39.
12. **Close B, Banister K, Baumans V, Bernoth EM, Bromage N, Bunyan J, Erhardt W, Flecknell P, Hackbarth NGH, Morton D, Warwick C.** 1997. Recommendation for euthanasia of experimental animals: part 2. *Lab Anim* **31**:14–15.
13. **Coenen AM, Drinkenburg WM, Hoenderken R, van Luijtelaar EM.** 1995. Carbon dioxide euthanasia in rats: oxygen supplementation minimizes signs of agitation and asphyxia. *Lab Anim* **29**:262–268.
14. **Colburn, D.** Personal communication.

15. **Committee on Guidelines for the Use of Animals in Neuroscience and Behavioral Research.** 2003. Guidelines for the Care and Use of Mammals in Neuroscience and Behavioral Research. Washington (DC): National Academies Press. p 102–108.
16. **Conahan ST, Narayan S, Vogel WH.** 1985. Effect of decapitation and stress on some plasma electrolyte levels in rats. *Pharmacol Biochem Behav* **23**:147–149.
17. **Coppola DM, Millar LC, Chen CJ, Vandenbergh JG.** 1997. Chronic cocaine exposure affects stimulus-induced but not spontaneous behavior of the near-term mouse fetus. *Pharmacol Biochem Behav* **58**:793–799.
18. **Cordo MG, Biggio G, Gessa GL.** 1980. Brain nucleotides in naïve and handling-habituated rats: differences in levels and drug sensitivity. *Brain Res* **188**:287–290.
19. **Danneman PJ, Mandrell TD.** 1997. Evaluation of five agents/methods for anesthesia of neonatal rats. *Lab Anim Sci* **47**:386–395.
20. **Danneman PJ, Stein S, Walshaw SO.** 1997. Humane and practical implications of using carbon dioxide mixed with oxygen for anesthesia or euthanasia of rats. *Lab Anim Sci* **47**:376–385.
21. **Depocas F, Behrens WA.** 1977. Effects of handling, decapitation, anesthesia and surgery on plasma noradrenaline levels in the white rat. *Can J Physiol Pharmacol* **55**:212–219.
22. **Dunn J, Scheving L.** 1971. Plasma corticosterone levels in rats killed sequentially at the ‘trough’ or ‘peak’ of the adrenocortical cycle. *J Endocrinol* **49**:347–348.
23. **Engel SR, Gaudet EA, Jackson KA, Allan AM.** 1966. Effect of in vivo administration of anesthetics on GABA_A receptor function. *Lab Anim Sci* **46**:425–429.
24. **Essential principles and practices.** 2003. Handbook of laboratory animal science, 2nd ed, vol 1. In: Hau J, Van Hoosier G Jr, editors. Boca Raton (FL): CRC Press.
25. **Feldman DB, Gupta BN.** 1976. Histopathological changes in laboratory animals resulting from various methods of euthanasia. *Lab Anim Sci* **26**:218–221.
26. **Fitzgerald M, Beggs S.** 2001. The neurobiology of pain: developmental aspects. *Neuroscientist* **7**:246–257.
27. **Fomby LM, Wheat TM, Hartter DE, Tuttle RL, Black CA.** 2004. Use of CO₂/O₂ anesthesia in the collection of samples for serum corticosterone analysis from Fischer 344 rats. *Contemp Top Lab Anim Sci* **43**:8–12.
28. **Gupta A, Cheng J, Wang S, Barr GA.** 2001. Analgesic efficacy of ketorolac and morphine in neonatal rat pups. *Pharmacol Biochem Behav* **68**:635–640.
29. **Hewett TA, Kovacs MS, Artwohl JE, Bennett BT.** 1993. A comparison of euthanasia methods in rats, using carbon dioxide in prefilled and fixed flow rate filled chambers. *Lab Anim Sci* **43**:579–582.
30. **Himwich WA.** 1962. Biochemical and neurophysiological development of the brain in the neonatal period. *Int Rev Neurobiol* **4**:117–1159.
31. **Hornett TD, Haynes AP.** 1984. Comparison of carbon dioxide/air mixture and nitrogen/air mixture for the euthanasia of rodents, design of a system for inhalation euthanasia. *Anim Technol* **35**:93–99.
32. **Howard HL, McLaughlin-Taylor E, Hill RL.** 1990. The effect of mouse euthanasia technique on subsequent lymphocyte proliferation and cell mediated lympholysis assays. *Lab Anim Sci* **40**:510–514.
33. **Iwarsson K, Rehinder C.** 1993. A study of different euthanasia techniques in guinea pigs, rats and mice. Animal response and post-mortem findings. *Scan J Lab Anim Sci* **20**:191–205.
34. **Jones DM, Arters J, Berger-Sweeney J.** 1999. Carbon dioxide-induced anaesthesia has no effect on brain biogenic amine concentration in mice. *Lab Anim Sci* **49**:316–318.
35. **Kasten T, Colliver JA, Montrey RD, Dunaway GA.** 1990. The effects of various anesthetics on tissue levels of fructose-2,6-bisphosphate in rats. *Lab Anim Sci* **40**:399–401.
36. **Kaufman W.** 2000. In: Krinke GJ, editor. The laboratory rat. San Diego (CA): Academic Press. p 227–242.
37. **Klaunberg BA, O'Malley J, Clark T, Davis JA.** 2004. Euthanasia of mouse fetuses and neonates. *Contemp Top Lab Anim Sci* **43**(5):29–34.
38. **Knott PJ, Hutson PH, Curzon G.** 1977. Fatty acid tryptophan changes on disturbing groups of rats and caging them single. *Pharmacol Biochem Behav* **7**:245–252.
39. **Leach MC, Howell VA, Allan TF, Morton DB.** 2002. Aversion to gaseous euthanasia agents in rats and mice. *Comp Med* **52**:249–257.
40. **Leach MC, Howell VA, Allan TF, Morton DB.** 2002. Degrees of aversion shown by rats and mice to different concentrations of inhalational anaesthetics. *Vet Record* **150**:808–815.
41. **Mathe AA, Stenfors C, Brodin E, Theodorsson E.** 1990. Neuropeptides in brain: effects of microwave irradiation and decapitation. *Life Sciences* **46**:287–293.
42. **Mazurkiewica-Kwilecki.** 1980. Single and repeated air blast stress and brain histamine. *Pharmacol Biochem Beh* **12**:35–39.
43. **Mellor DJ, Gregory NG.** 2003. Responsiveness, behavioral arousal and awareness in fetal and newborn lambs: experimental, practical and therapeutic implications. *NZ Vet J* **51**:2–13.
44. **Methods of Behavioral Analysis in Neuroscience.** J. J. Buccafusco, editor. 2001. CRC Press.
45. **Miller JM, Jope RS, Ferraro TN, Hare TA.** 1990. Brain amino acid concentrations in rats killed by decapitation and microwave irradiation. *J Neurosci Methods* **31**:187–92.
46. **Nahas K, Provost J-P.** 2002. Blood sampling in the rat: current practices and limitations. *Comp Clin Path* **11**:14–37.
47. **National Institutes of Health Animal Research Advisory Committee.** 2004. Guidelines for the euthanasia of mouse and rat fetuses and neonates. [<http://oacu.od.nih.gov/ARAC/euthmous.pdf>].
48. **National Research Council.** 1996. Guide for the care and use of laboratory animals. p 65–66.
49. **Nazian SJ.** 1988. Serum concentrations of reproductive hormones after administration of various anesthetics to immature and young adult male rats. *Proceedings of the Society for Experimental Biology and Medicine* **187**:482–487.
50. **O'Connor JL, Kellom TA.** 1989. Ether as an anesthetic for decapitation in the rat: gonadotropin secretion by subsequently established anterior pituitary cell cultures (42866). *Proceedings of the Society for Experimental Biology and Medicine* **190**:320–323.
51. **Office of Laboratory Animal Welfare, National Institutes of Health, U.S. Department of Health and Human Services.** 2002. Public Health Service Policy on Humane Care and Use of Laboratory Animals, Clarification Regarding Use of Carbon Dioxide for Euthanasia of Small Laboratory Animals. [<http://grants.nih.gov/grants/guide/notice-files/NOT-OD-02-062.html>].
52. **Pecaut MJ, Smith AL, Jones TA, Gridley DS.** 2000. Modification of immunologic and hematologic variables by method of CO₂ euthanasia. *Comp Med* **50**:595–602.
53. **Pettinger WA, Tanaka K, Keeton K, Campbell WB, Brooks SN.** 1975. Renin release, an artifact of anesthesia and its implications in rats. *Proceedings of the Society for Experimental Biology and Medicine* **148**:625–630.
54. **Phifer CB, Terry LM.** 1986. Use of hypothermia for general anesthesia in preweanling rodents. *Physiol Behav* **38**:887–890.
55. **Pritchett KR, Corrow D, Stockwell JD, Smith AL.** 2005. Euthanasia of neonatal mice using carbon dioxide. *Comp Med* **55**:275–281.
56. **Reilly JS, Rose MA.** 2001. Scientific considerations—influence of methods of euthanasia on scientific data. *Euthanasia of Animals Used for Scientific Purposes.* ANZCCART, 17–24.
57. **Robinson SE, Wallace MJ.** 2001. Effect of perinatal buprenorphine exposure on development in the rat. *J Pharmacol Exp Ther* **298**:797–804.
58. **Ronca AE, Alberts JR.** 1995. Simulated uterine contractions facilitate fetal and newborn respiratory behavior in rats. *Physiol Behav* **5**:1035–41.
59. **Schnecko AK, Witte K, Lemmer B.** 1998. Effects of routine procedures on cardiovascular parameters of Sprague-Dawley rats in periods of activity and rest. *J Exp An Sci* **38**:181–190.
60. **Schriefer JA, Plunkett WC, Hassen AH.** 1989. Decapitation increases plasma sodium and potassium in the rat. *J Pharmacol Methods* **21**:155–159.
61. **Segel LD, Rendig SV.** 1986. Sodium pentobarbital effects on cardiac function and response to dobutamine. *J Cardiovasc Pharmacol* **8**:392–397.

62. **Sharp J, Zammit T, Azar T, Lawson D.** 2003. Stress-like responses to common procedures in individually and group-housed female rats. *Contemp Top Lab Anim Sci* **42**:9–18.
63. **Singer D.** 1999. Neonatal tolerance to hypoxia: a comparative-physiological approach. *Comp Biochem Physiol* **123**:221–234.
64. **Slott VL, Linder DE, Dyer CJ.** 1994. Method of euthanasia does not affect sperm motility in the laboratory rat. *Reprod Toxicology* **8**:317–74.
65. **Smotherman WP, Robinson SR.** 1985. The rat fetus in its environment: behavioral adjustments to novel, familiar, aversive, and conditioned stimuli present *in utero*. *Behav Neurosci* **99**:521–530.
66. **Urbanski HF, Kelley ST.** 1991. Sedation by exposure to a gaseous carbon dioxide-oxygen mixture: application to studies involving small laboratory species. *Lab Anim Sci* **41**:80–82.
67. **Vannucci RC, Wolf JW.** 1977. Oxidative metabolism in fetal rat brain during maternal halothane anesthesia. *Environ Health Perspect* **21**:215–219.
68. **Varki AP, Fritz JL, Davis RB.** 1979. Effects of cervical dislocation on colony-forming cells in murine marrow cultures. *Exp Hematology* **7**:397–400.
69. **Walter G.** 2000. Effects of carbon dioxide inhalation on hematology, coagulation, and serum clinical chemistry values in rats. *Toxicol Path* **27**:217–225.
70. **Woodbury CJ, Ritter AM, Koerber HR.** 2001. Central anatomy of individual rapidly adapting low-threshold mechanoreceptors innervating the “hairy” skin of newborn mice: early maturation of hair follicle afferents. *J Comp Neurol* **436**:304–323.
71. **Wuttke W, Meites J.** 1970. Effects of ether and pentobarbital on serum prolactin and LH levels in proestrous rats. *Proc Soc Exp Biol Med* **135**:648–652.
72. **Yamamoto Y, Hasegawa H, Ikeda K, Ichlyama A.** 1988. Cervical dislocation of mice induces rapid accumulation of platelet serotonin in the lung. *Agents and Actions* **25(1/2)**:49–56.
73. **Yi DK, Barr GA.** 1997. Formalin-induced c-fos expression in the spinal cord of fetal rats. *Pain* **73**:347–354.
74. **Zarembka FR, Koller DE, Plotka ED.** 1989. Effect of ether or ketamine anesthesia on rat uterine estrogen and progesterone receptors. *Clin Chem* **35**:143–145.